

**Clerk of the Circuit Court
2022 Judicial Statistics**

	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	YTD
AD Adoption	1	2	3	2	0	0							8
CA Court Administration	1	0	0	0	0	0							1
CC Contempt of Court	2	1	0	4	0	2							9
CF Criminal Felony	52	29	45	25	43	57							251
CH Chancery	1	1	1	0	3	5							11
CL Civil Law Violation	1	0	1	0	0	0							2
CM Criminal Misdemeanor	17	25	26	20	24	37							149
CV Conservation Violation	1	0	1	0	2	2							6
DC Dissolution with Children	22	16	20	24	17	21							120
DN Dissolution without Children	13	6	18	16	15	14							82
DT DUI	24	22	20	21	24	26							137
DV Domestic Violence	18	13	13	14	8	8							74
ED Eminent Domain	0	0	0	4	12	3							19
EV Eviction	24	18	31	47	29	28							177
FA Family	12	24	28	21	26	13							124
FC Foreclosures	12	18	31	18	32	20							131
GR Guardianships	9	2	6	11	4	7							39
JV Juvenile	0	0	0	0	4	2							6
JA Juvenile Abuse/Neglect	5	6	6	3	9	11							40
JD Juvenile Delinquency	8	19	23	7	5	21							83
LA Law	6	13	11	8	12	5							55
LM Law Magistrate	24	11	36	10	14	31							126
MH Mental Health	2	1	0	0	1	0							4
MR Misc. Remedy	7	6	9	6	8	5							41
MT Major Traffic	107	106	113	95	120	125							666
MX Misc Criminal	20	9	21	12	22	15							99
OP Order of Protection	20	33	33	37	38	41							202
OV Ordinance Violation	2	0	2	3	4	4							15
PR Probate	13	6	16	6	8	17							66
QC Quasi Criminal	0	0	0	0	0	0							0
SC Small Claims	114	63	124	84	92	101							578
TR Traffic	280	288	357	420	299	301							1945
TX Tax	6	6	1	9	1	5							28
WI Wills	22	19	10	16	17	19							103
XX Misc	2	1	0	0	3	3							9
	848	764	1006	943	896	949	0	0	0	0	0	0	5406
Totals for 2021	683	675	949	893	749	956	801	965	859	812	871	852	10065



KENDALL COUNTY HEALTH DEPARTMENT

Are you or someone you love struggling with opioid use?

We are here to help.

Kendall County Health Department is making Naloxone available* for free to Kendall County community members Naloxone can be picked up the Health Department, located at 811 W. John St., Yorkville, IL. For questions we can be reached at (630) 553-9100.

What is Naloxone?

Naloxone (also known as Narcan®) is a medication that can reverse an overdose caused by an opioid drug (heroin or prescription pain medications). When administered during an overdose, naloxone blocks the effects of opioids on the brain and quickly restores breathing. Naloxone has one critical function: to reverse the effects of opioids in order to prevent overdose death. Naloxone has no potential for abuse. Naloxone DOES NOT reverse overdoses that are caused by non-opioid drugs, such as cocaine, benzodiazepines (e.g. Xanax, Klonopin and Valium), methamphetamines, or alcohol.

When to give naloxone?

You suspect an opioid overdose if you see:

- Very sleepy or unresponsive
- Breathing is slow or stopped
- Pupils are tiny
- Lips or nails are blue or gray
- Skin feels cold or clammy

How to administer naloxone?

For the use of naloxone, become familiar with and carefully follow the included manufacturer's product instructions. A quick reference guide can be found on the reverse side of this page.

To learn more about our mental health and substance misuse programs and services please feel free to contact us at (630) 553-9100 or visit us at www.kendallhealth.org

*One kit per requestor, comprised of two doses of Narcan® nasal spray; and while quantities last



NARCAN[®] (naloxone HCl) **NASAL SPRAY**

QUICK START GUIDE Opioid Overdose Response Instructions

Use NARCAN Nasal Spray (naloxone hydrochloride) for known or suspected opioid overdose in adults and children.

Important: For use in the nose only.

Do not remove or test the NARCAN Nasal Spray until ready to use.

1 Identify Opioid Overdose and Check for Response

Ask person if he or she is okay and shout name.

Shake shoulders and firmly rub the middle of their chest.

Check for signs of opioid overdose:

- Will not wake up or respond to your voice or touch
 - Breathing is very slow, irregular, or has stopped
 - Center part of their eye is very small, sometimes called “pinpoint pupils”
- Lay the person on their back to receive a dose of NARCAN Nasal Spray.**



2 Give NARCAN Nasal Spray

Remove NARCAN Nasal Spray from the box.

Peel back the tab with the circle to open the NARCAN Nasal Spray.



Hold the NARCAN nasal spray with your thumb on the bottom of the plunger and your first and middle fingers on either side of the nozzle.




Gently insert the tip of the nozzle into either nostril.

- Tilt the person's head back and provide support under the neck with your hand. Gently insert the tip of the nozzle into **one nostril**, until your fingers on either side of the nozzle are against the bottom of the person's nose.



Press the plunger firmly to give the dose of NARCAN Nasal Spray.

- Remove the NARCAN Nasal Spray from the nostril after giving the dose.



Get emergency medical help right away.

Move the person on their side (recovery position) after giving NARCAN Nasal Spray.

Watch the person closely.

If the person does not respond by waking up, to voice or touch, or breathing normally another dose may be given. NARCAN Nasal Spray may be dosed every 2 to 3 minutes, if available.



3 Call for emergency medical help, Evaluate, and Support

Repeat Step 2 using a new NARCAN Nasal Spray to give another dose in the other nostril. If additional NARCAN Nasal Sprays are available, repeat step 2 every 2 to 3 minutes until the person responds or emergency medical help is received.

For more information about NARCAN Nasal Spray, go to www.narcannasalspray.com, or call 1-844-4NARCAN (1-844-462-7226).